

the elytra are oval, broad, truncated in front, convex, striated; the first stria bi-furcated in front; the margin impressed; legs and antennæ of the general colour; mandibulæ and labrum dark; palpi of a light yellowish brown.

Adelaide.

*Harpalus Boisduvalii*: length  $3\frac{3}{4}$ '; very nearly allied to *Australis*, Dej., but having the thorax broader, shorter, more rounded laterally; the elytra broader and not so deeply striated; the thorax is often of a brilliant and rather metallic colour.

Swan River.

*Harpalus Bostockii*: length 4'; of a brilliant green copper colour; head rather small, rounded; thorax transverse, about as broad behind as in front, rounded laterally, with the posterior angles obtuse, and the anterior rounded; the impressions are moderate; elytra broad, rather depressed, marked with slight striæ, the intervals of which are plane; the sutural stria deviates towards the scutellum, and there is a light abbreviate stria; a small punctiform impression is seen on the posterior part between the second and third striæ; the margin is impressed; legs of a brownish red, with the thighs and end of the tibiæ black; antennæ dark, with the two first articles red, as are the palpi.

Swan River.

*Harpalus Versicolor*: length 4'; broad; of a reddish brown; elytra of a dark brown; head almost round, rather rugous, and impressed in front; thorax transverse, rather cordiform, much narrower behind than in front, with a light longitudinal sulcate, and two equally light posterior impressions; elytra strongly striated, almost sulcated, impressed laterally; no abbreviate stria near the scutellum; the humeral part of the margin is yellow; lower side of the body of a light brown; legs and parts of the mouth yellow.

Swan River.

*Harpalus Deyrolii*: length  $3\frac{1}{2}$ '; of a light bronzed colour; head rounded, with a transversal line in front; thorax transverse, rounded laterally, as broad behind as in front; having a well marked transverse impression in front, a feeble longitudinal sulcate and two weak impressions behind; elytra oval, with feeble longitudinal striæ, rather far apart one from the other, and the intervals of which are plane; the sutural one deviates towards the scutellum, and there is a very short abbreviate stria; margin very feebly impressed; lower

side of the body of a light brown; legs, parts of the mouth and antennæ yellow.

Port Lincoln, South Australia.

*Harpalus Femoralis*: length  $3\frac{1}{4}$ '; black; rather brilliant; head with two punctiform impressions united by a transversal line in front; thorax almost square, transverse; as broad behind as in front, having all its impressions feebly marked; elytra rather broad, strongly striated; the second stria bi-furcated near the scutellum; posterior part of the margin impressed; a feebly marked punctiform impression behind on the interval between the third and fourth striæ; thighs black; tibiæ, tarsi, palpi and antennæ of a light red.

Rockhampton, Queensland.

*Harpalus Mandibularis*: length  $2\frac{3}{4}$ '; of a bronzed green colour; body rather broad and depressed; head smooth, with a strong sulcate in front; thorax broad, transversal, with the sides very much rounded but straight behind; the anterior angles are rounded, and the posterior straight; the central longitudinal sulcate and the two posterior impressions are strong, the anterior transverse one is feeble; elytra rather broader than the thorax, oval, depressed, having longitudinal striæ, the intervals of which are not convex; a very feeble abbreviate stria near the scutellum; a row of impressions on the margin; lower side of the body brown; legs, palpi, and base of the antennæ yellow; remainder of the latter a little darker; mandibulæ yellow at the base, and black towards their extremity.

Port Denison.

*Harpalus Dingo*: length  $2\frac{1}{2}$ ' to 3'; general form rather depressed; of a dark brilliant brown; head large, rounded, having a faint impression in front; thorax almost square, broader in front than behind, rounded laterally; the impressions very faint; scutellum brown; elytra oval, striated; an abbreviate stria near the scutellum; a very faint punctiform impression behind, between the second and third striæ; margin having on its posterior half a series of strong impressions; lower side of the body black; inferior margin of the elytra brown; legs, palpi, and three first articles of the antennæ yellow, the remaining of the latter obscure.

Rockhampton.

*Harpalus Novæ Hollandiæ*: length  $3\frac{1}{2}$ '; nearly allied to *Rubripes*, but smaller; of an æneus copper green; head with two impressions before the eyes; thorax rather transversal, rounded and bordered laterally; narrower in front