

141.—*BEMBIDIUM BIPUSTULATUM*. n. sp.

Length $1\frac{1}{4}$ lines.

Black, subnitid. Head smooth and longitudinally impressed on each side between the eyes. Thorax rather flat, much broader than the length, slightly rounded on the sides, not narrowed behind, and truncate at the base, with the median line scarcely visible and the basal impressions very short. Elytra broad, striated and of a bronzy hue, with an indistinct red spot near the apex on the outside of each elytron. Legs, palpi, and antennæ yellow.

142.—*BEMBIDIUM PUNCTIPENNE*. n. sp.

Length 1 line.

Entirely of a glossy red. Head large and flat, with the impressions between the eyes long and deep. Thorax transversal, convex, much rounded on the sides and narrowed at the base, with the median line lightly marked. Elytra moderately convex, and marked with regular rows of distinct punctures which disappear towards the apex. The suture and lateral margins of the elytra are of a rather darker hue than the rest of the body.

143.—*BEMBIDIUM ATRICEPS*. n. sp.

Length 1 line.

Head black, and largely impressed on each side between the eyes. Thorax yellowish red, rather flat, transversal, slightly rounded on the sides anteriorly, and with the posterior angles square and broadly reflexed. Elytra somewhat flat, strongly striated, punctured in the striæ, and of a reddish colour, with the suture, apex, and a broad median fascia, black. Legs, antennæ, and palpi yellow.

144.—*BEMBIDIUM TRANSVERSICOLLE*. n. sp.

Length $1\frac{1}{6}$ lines.

Of a pale red, slightly clouded with brown on the head and elytra. Thorax transversal, convex, rounded on the sides in front, square and acute at the posterior angles, and transversely de-